


PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 31/00	A2	(11) International Publication Number: WO 00/40226 (43) International Publication Date: 13 July 2000 (13.07.00)
(21) International Application Number: PCT/US99/27951 (22) International Filing Date: 20 December 1999 (20.12.99) (30) Priority Data: 60/114,840 6 January 1999 (06.01.99) US 60/115,051 8 January 1999 (08.01.99) US 60/115,922 14 January 1999 (14.01.99) US 60/120,543 17 February 1999 (17.02.99) US (71) Applicant (for all designated States except US): PHARMACIA & UPJOHN COMPANY [US/US]; 301 Henrietta Street, Kalamazoo, MI 49001 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): MEGLASSON, Martin, Durham [US/US]; 5337 Whippoorwill Drive, Kalamazoo, MI 49002 (US). McCALL, Robert, B. [US/US]; 413 Sunview, Kalamazoo, MI 49002 (US). (74) Agent: STEIN, Bruce; Intellectual Property Legal Services, Pharmacia & Upjohn Company, 301 Henrietta Street, Kalamazoo, MI 49001 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: METHOD OF TREATING SEXUAL DISTURBANCES		
(57) Abstract		
<p>The present invention is a method of treating sexual disturbances in humans and inducing mating in non-human mammals using the compounds of formula (A) in a dosage range where the sexually therapeutic amount is from about 0.2 thru 8 mg/person/dose and where the sexually mating amount is from about 0.003 thru 0.2 mg/kg/dose.</p> <div data-bbox="852 1165 1323 1417"><p style="text-align: right;">(A)</p></div> <div data-bbox="958 1795 1396 1932"><p>Atty. Docket No. 6794S-000028/US/01 Serial No. 10/678,212 Michael Hawley et al Reference 44</p></div>		

EV 726254284 US

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

METHOD OF TREATING SEXUAL DISTURBANCES1. Field of the Invention

The present invention is a method of treating sexual disturbances using the
5 compounds of US Patent 5,273,975.

2. Description of the Related Art

There are a number of diseases/conditions which directly affect the sexual lives of humans and animals. These include orgasmic disorders, lack of interest in sex and erectile dysfunction in males. In addition there are a number of diseases/conditions which
10 indirectly affect the sexual lives of humans. There are a few pharmaceutical agents to treat this diseases/conditions and others in clinical development.

US Patent 5,273,975 discloses that the compounds (A) of the present invention are useful to stimulate sexual activity and to alleviate impotence. The useful dosage range disclosed in US Patent 5,273,975 is "at least 10 mg up to about 1200 mg per day". The
15 operable dosage range in the present invention is considerably lower. The compounds of US Patent 5,273,975 have been used in clinical trials to prove their usefulness in treating Parkinson's disease at a dose of 30 mg/person. Compounds (A) of the present invention are used at a dose of less than 8 mg in treating the sexual disturbances of the present invention. US Patent 5,273,975 generically discloses compounds where the variable substituent "A"
20 can be either a carbonyl group (-CO-) or a thiocarbonyl (-CS-).

US Patent 5,250,534 discloses sildenafil (VIAGRA®). The compounds (A) of the present invention are quite different chemically than sildenafil. International Publication WO94/282902 discloses the use of sildenafil for treating male impotence.

US Patent 4,127,118 discloses the intracavernosal injection of a vasodilator for
25 enhancing an erection of the penis. PGE1 is a vasodilator and is sold as CAVERJECT® to treat male erectile dysfunction. The compounds (A) of the present invention are not prostaglandins and are not injected into the penis.

International Journal of Impotence Research, 10 (Supplement 3), August 1998, Abstracts #417 and 419 disclose that apomorphine, which is available in a number of
30 countries to treat Parkinson's Disease, is in late-stage development as a sublingual formulation for treating male erectile dysfunction. The compounds (A) used in the present invention are not apomorphine analogs. US Patent 5,770,606 discloses sublingual use of apomorphine for treating erectile dysfunction.

US Patent 5,541,211 discloses that yohimbine can be used to treat erectile dysfunction. The compounds of formula (A) are quite different chemically as compared to yohimbine.


US Patent 4,801,587 discloses that phentolamine (VASOMAX®) which is available in a number of countries for treating hypertension is also useful for treating erectile dysfunction. The compounds of formula (A) are quite different chemically as compared to phentolamine. In addition, the compounds of formula (A) are not applied either topically to the penis nor intra-urethrally.

US Patent 5,773,020 discloses that intraurethral PGE1 (MUSE) is useful for treating erectile dysfunction. Compounds (A) are not prostaglandins and are not administered intraurethrally.

Brain Research, 55, 383-389 (1973) reports on the sexually stimulating effect of L-DOPA given to male rats and conjectures that it may be the reason that hypersexuality occasionally is seen in patients with Parkinson's syndrome during L-DOPA treatment.

SUMMARY OF INVENTION

Disclosed is a method of treating sexual disturbances in a human who is in need of such treatment which comprises administering a sexually therapeutically effective amount of a compound of the formula (A)



where

R₁, R₂ and R₃ are the same or different and are: -H, C₁-C₆ alkyl, C₃-C₅ alkenyl, C₃-C₅ alkynyl, C₃-C₅ cycloalkyl, C₄-C₁₀ cycloalkyl, phenyl substituted C₁-C₆ alkyl, -NR₁R₂ where R₁ and R₂ are cyclized with the attached nitrogen atom to produce pyrrolidyl, piperidyl, morpholinyl, 4-methyl piperazinyl or imidazolyl;

X is: -H, C₁-C₆ alkyl, -F, -Cl, -Br, -I, -OH, C₁-C₆ alkoxy, cyano, carboxamide, carboxyl, (C₁-C₆ alkoxy)carbonyl;

A is: CH, CH₂, CH-(halogen) where halogen is -F, -Cl, -Br, -I, CHCH₃, C=O, C=S, C-SCH₃, C=NH, C-NH₂, C-NHCH₃, C-NHCOOCH₃, C-NHCN, SO₂, N;

B is: CH₂, CH, CH-(halogen) where halogen is as defined above, C=O, N, NH,

N-CH₃,

D is: CH, CH₂, CH-(halogen) where halogen is as defined above, C=O, O, N, NH, N-CH₃; and n is 0 or 1, and where — is a single or double bond, with the provisos:

(1) that when n is 0, and

5 A is CH₂, CH-(halogen) where halogen is as defined above, CHCH₃, C=O, C=S, C=NH, SO₂; then

D is CH₂, CH-(halogen) where halogen is as defined above, C=O, O, NH, N-CH₃;

(2) that when n is 0, and

10 A is CH, C-SCH₃, C-NH₂, C-NHCH₃, C-NHCOOCH₃, C-NHCN, N; then

D is CH, N;

(3) that when n is 1, and

A is CH₂, CH-(halogen) where halogen is as defined above, CHCH₃, C=O, C=S, C=NH, SO₂; and

15 B is CH₂, CH-(halogen) where halogen is as defined above, C=O, NH, N-CH₃; then

D is CH₂, C=O, O, NH, N-CH₃;

(4) that when n is 1, and

A is CH, C-SCH₃, C-NH₂, C-NHCH₃, C-NHCOOCH₃, C-NHCN, N; and

20 B is CH, N; then

D is CH₂, C=O, O, NH, N-CH₃;

(5) that when n is 1, and

A is CH₂, CHCH₃, C=O, C=S, C=NH, SO₂, and

B is CH, N; then

25 D is CH, N; and pharmaceutically acceptable salts thereof to the human.

Also disclosed is a method of inducing mating a non-human mammal which comprises administering a sexually mating amount of a compound of the formula (A) and pharmaceutically acceptable salts thereof.

Further disclosed is a method of treating a sexual deficiency state in a human who
30 has epilepsy, craniopharyngioma, hypogonadism or who has had a hysterectomy/oophorectomy, hysterectomy or oophorectomy which comprises administering a sexually therapeutically effective amount of a compound of the formula (A) and pharmaceutically acceptable salts thereof.

Additionally disclosed is a method of increasing sexual desire, interest or performance in a human who is desirous thereof which comprises administering a sexually useful effective amount of a compound of the formula (A) and pharmaceutically acceptable salts thereof.

- 5 Disclosed is (5R)-5-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinoline-2(1H)-thione and pharmaceutically acceptable salts thereof.

DETAILED DESCRIPTION OF THE INVENTION

The present invention includes four separate methods of treating sexual problems. First, it is a method of treating sexual disturbances in a human who is in need of such
10 treatment which comprises administering a sexually therapeutic amount of a compound of the formula (A) and its pharmaceutically acceptable salts. Second, it includes a method of inducing mating a non-human mammal which comprises administering a sexually mating amount of a compound of the formula (A) and pharmaceutically acceptable salts thereof. Thirdly, it includes a method of treating a sexual deficiency state in a human who has a
15 disease/condition the primary pathology of which does not relate to sexual dysfunction, but indirectly produces reduced sexual functioning. The diseases/conditions which can produce this type of sexual dysfunction include epilepsy, craniopharyngioma, hypogonadism or who has had a hysterectomy/oophorectomy, hysterectomy or oophorectomy. Fourthly, there are individuals who do not appear to have any sexual disturbance or disease/condition which
20 will produced an adverse effect on their sexual lives, but none the less wish to increase their sexual desire, interest or sexual performance and who will benefit by use of the compounds of formula (A).

The compounds of formula (A), and pharmaceutically acceptable salts, which are useful in the method of treatment of the present invention are known, see US Patent
25 5,273,975. It is preferred that the pharmaceutically acceptable salt is selected from the group consisting of salts of the following acids methanesulfonic, hydrochloric, hydrobromic, sulfuric, phosphoric, nitric, benzoic, citric, tartaric, fumaric, maleic, $\text{CH}_3-(\text{CH}_2)_n-\text{COOH}$ where n is 0 thru 4, $\text{HOOC}-(\text{CH}_2)_n-\text{COOH}$ where n is as defined above. It is preferred that the compound of formula (A) have n be 0, A be C=O and D be NH. It is more preferred
30 that the compound of formula (A) be either (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one or (5R)-5-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinoline-2(1H)-thione. It is more preferred that the compound of formula (A) be (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (Z)-2-butenedioate (1:1).

A preferred process to produce the preferred compounds within the scope of the compounds of formula (A) is set forth in PREPARATION 1 and the numerical EXAMPLEs as well as CHART A.

Non-human mammals include commercial animals (horses, cattle, swine, sheëp and
5 transgenic mice) and zoo animals (panda bears, elephants, zebras, lions, tigers, monkeys and apes), sporting animals (horses and dogs) as well as domesticated animals (dogs and cats). The humans and non-human mammals both be either a male or female.

The sexual disturbances in humans treated by the present invention include hypoactive sexual desire disorder, female sexual arousal disorder, male erectile disorder,
10 female orgasmic disorder and male orgasmic disorder. The sexual disturbances of the present invention are known to those skilled in the art and are adequately described for a medical practitioner. In particular, see Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, (DSM-IV), American Psychiatric Association, Washington DC, 1994 as well as the DSM-IV Guidebook, American Psychiatric Press, Inc., Washington DC,
15 1995.

When inducing mating in the non-human mammals it is desirable to treat one member of the mating pair in some cases. This is most likely to be the situation where previous attempts at mating the pair have failed and one member of the pair is not known to have prior offspring. In that case, this member would be advantageously treated. In other
20 cases, it may be desirable to treat both the male and female prior to the anticipated mating. The latter is most likely to be advantageous when the intended mating is between sexually inexperienced animals and prior attempts at mating the pair have failed.

The compounds of formula (A) are also useful in treating humans whose primary disease, condition or situation is not one of sexual dysfunction, but rather the primary
25 disease or condition leads to a secondary clinical situation where the human (male or female) experiences diminished sexual desire, interest and/or function. These diseases include epilepsy, craniopharyngioma, hypogonadism or who has had a hysterectomy/oophorectomy, hysterectomy or oophorectomy and can produce a sexual deficiency state which is treated in the same way as those individuals who have one or more
30 of the five sexual disturbances identified above. The sexually therapeutic effective amount is the same and is administered in the same way by the same routes of administration as for those humans who have sexual disturbances.

The compounds of formula (A) are also useful in increasing sexual desire, interest or performance in individuals who do not have any of the five sexual disturbances nor any

of the following diseases or conditions, epilepsy, craniopharyngioma, hypogonadism or who has had a hysterectomy/oophorectomy, hysterectomy or oophorectomy. These individuals are treated the same way as those having any of the five sexual disturbances. The sexually useful effective amount to be used here is the same as the sexually therapeutically effective amount for those with sexual disturbances. The sexually useful therapeutically effective amount is administered in the same way by the same routes of administration as for those humans who have sexual disturbances.

In (1) treating the sexual disturbances of humans, (2) inducing mating of non-human mammals, (3) treating humans whose primary disease or condition leads to a secondary clinical situation where the human experiences diminished sexual desire, interest and/or function and (4) increasing the sexual desire, interest or performance in individuals who do not have any of the above problems the compounds of formula (A) in the known appropriate pharmaceutical dosage form for a given route of administration may be administered orally, intra-nasally, buccally, intra-pulmonary, parenterally and rectally. For treating humans, it is preferred that the compound of formula (A) be administered orally, intra-nasally, buccally and intra-pulmonarily; it is more preferred that the compounds of formula (A) be administered orally. For treating non-human mammals, it is preferred that the compounds of formula (A) be administered orally, parenterally and rectally; it is more preferred that the compounds of formula (A) be administered orally.

The operable sexually therapeutic effective amount as well as the operable sexually useful effective amount of the compounds of formula (A) is from about 0.2 thru about 8 mg/person/dose. It is preferred that the sexually therapeutic effective amount and sexually useful effective amount is from about 0.5 thru about 5 mg/person/dose. It is more preferred that the sexually therapeutic effective amount and sexually useful effective amount is from about 1 thru about 3 mg/person/dose. If doses less than this are used the desired effect will not be obtained. If doses greater than this are used, undesirable side effects occur and the desired effect will not be obtained. The term "therapeutic" is used in treating a disease or condition; the term "useful" is used when treating humans who do not have one of the above diseases or conditions which cause sexual dysfunction but who are considered normal and who wish to increase their sexual desire, interest or performance.

The operable sexually mating amount of the compounds of formula (A) is from about 0.003 thru about 0.2 mg/kg/dose. It is preferred that the sexually mating amount is from about 0.01 thru about 0.125 mg/kg/dose. It is more preferred that the sexually mating amount is from about 0.025 thru about 0.075 mg/kg/dose.

To obtain the desired effect in humans, the sexually therapeutic effective amount or sexually useful effective amount of the compounds of formula (A) should be administered from about 10 minutes to about 8 hr prior to sexual activity. It is preferred that the compounds of formula (A) be administered from about 0.5 hr to about 1 hr prior to sexual activity. It is more preferred that the compounds of formula (A) be administered about 0.5 hr prior to sexual activity. Sexual activity includes sexual intercourse with or without orgasm, ejaculation, masturbation and sexual foreplay.

To obtain the desired mating effect in non-human mammals, the compounds of formula (A) is administered from about 10 minutes to about 8 hr prior to mating. It is preferred that the compounds of formula (A) be administered from about 10 minutes to about 1 hr prior to mating. It is more preferred that the compounds of formula (A) be administered from about 10 minutes to about 0.5 hr prior to mating.

It is preferred that the humans do not have Parkinson's disease.

It is preferred that the in humans administration of the compounds of formula (A) does not produce postural hypotension.

The present invention is further illustrated by EXAMPLES A-P.

The exact dosage of the compounds of formula (A) which are useful in the four methods of this invention depends on the route of administration, the particular compound used, the particular condition being treated, the severity of the condition being treated, the age, weight, general physical condition of the particular patient, other medication the individual may be taking as is well known to those skilled in the art and can be more accurately determined by measuring the blood level or concentration of the compound and/or metabolite in the patient's blood and/or the patient's response to the particular condition being treated.

The compounds of formula (A) can also be used with other agents in (1) treating the five sexual disturbances, (2) in inducing mating of a non-human mammal, (3) in treating a sexual deficiency state in humans who have one or more of the following diseases/conditions - epilepsy, craniopharyngioma, hypogonadism or who has had a hysterectomy/oophorectomy, hysterectomy or oophorectomy or (4) increasing sexual desire, interest or performance in a human who does not have any sexual disturbance or one of the above diseases/conditions. It is known that vascular smooth muscle relaxation is necessary for vascular engorgement of the penis and the clitoris during normal male and female sexual behavior and that agents which promote vascular engorgement of the penis and clitoris are useful in enhancing sexual motivation, desire and performance. Agents which

effectuate vascular smooth muscle relaxation include phosphodiesterase type 5 inhibitors, such as sildenafil (Viagra) and ICOS-351; phosphodiesterase type 3 inhibitors, such as milrinone; non-selective phosphodiesterase inhibitors such as papaverine; nitric oxide donor drugs such as linsidomine; alpha type 1 adrenergic receptor antagonists such as phentolamine produce relaxation of vascular smooth muscle from human penis in the presence of norepinephrine; alpha type 2 adrenergic receptor antagonists such as yohimbine; prostaglandin E1 receptor agonists such as PGE1 (CAVERJECT™ Injection and MUSE™); as well as vasoactive intestinal polypeptide (VIP) agents.

These vascular smooth muscle relaxation agents are administered by oral, parenteral, buccal, rectal, intranasal, intrapulmonary, intraurethral or topical routes. The preferred route of administration will depend on the specific properties of the drug(s) to be combined with the compound(s) of formula (A). The preferred route of administration for a phosphodiesterase type 3 or 5 receptor inhibitor and an alpha type 1 or 2 adrenergic receptor inhibitor is oral. The preferred route of administration for a prostaglandin E receptor agonist, nitric oxide donor, non-specific phosphodiesterase inhibitor, and VIP is intracavernosal injection or intraurethral topical administration in male mammals or topical administration to the vulva in the female.

The dose of some of the vascular smooth muscle relaxation agents is known to those skilled in the art, especially for sildenafil (VIAGRA™ Tablets), PGE1 (CAVERJECT™ Injection and MUSE™). For yohimbine the dose is about 1 to about 10 mg orally and can be given about three or four times daily as a routine. For phentolamine the dose is about 2 to about 40 mg orally or by buccal patch or about 0.5 to about 5 mg when given by intracavernosal injection. For papaverine the dose is about 4 to about 20 mg given by intracavernosal injection. For VIP the dose is about 5 to about 60 mg when given by intracavernosal injection.

When the compounds of formula (A) are used in conjunction with the vascular smooth muscle relaxation agents the compounds of formula (A) should be administered within the time frame discussed above. The sexually effective time period for administration of the vascular smooth muscle relaxation agents is well known for sildenafil (VIAGRA™ Tablets), PGE1 (CAVERJECT™ Injection and MUSE™). The sexually effective time period for administration of yohimbine is about 0 to about 4 hours before sexual activity; for phentolamine it is about 0 to about 4 hours prior to sexual activity when given orally or buccally and about 0 to about 2 hours when given by intracavernosal injection; for papaverine and VIP it is about 0 to about 2 hours prior to sexual activity.

When used in combination, the preferred combinations are (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one and sildenafil, (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one and PGE1, (5R)-5-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinoline-2(1H)-thione and sildenafil or
5 (5R)-5-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinoline-2(1H)-thione and PGE1.

DEFINITIONS AND CONVENTIONS

The definitions and explanations below are for the terms as used throughout this entire document including both the specification and the claims.

I. CONVENTIONS FOR FORMULAS AND DEFINITIONS OF VARIABLES

10 The carbon atom content of variable substituents is indicated in one of two ways. The first method uses a prefix to the entire name of the variable such as "C₁-C₄", where both "1" and "4" are integers representing the minimum and maximum number of carbon atoms in the variable. The prefix is separated from the variable by a space. For example, "C₁-C₄ alkyl" represents alkyl of 1 through 4 carbon atoms, (including isomeric forms
15 thereof unless an express indication to the contrary is given). Whenever this single prefix is given, the prefix indicates the entire carbon atom content of the variable being defined. Thus C₂-C₄ alkoxycarbonyl describes a group CH₃-(CH₂)_n-O-CO- where n is zero, one or two. By the second method the carbon atom content of only each portion of the definition is indicated separately by enclosing the "C_i-C_j" designation in parentheses and placing it
20 immediately (no intervening space) before the portion of the definition being defined. By this optional convention (C₁-C₃)alkoxycarbonyl has the same meaning as C₂-C₄ alkoxy-carbonyl because the "C₁-C₃" refers only to the carbon atom content of the alkoxy group. Similarly while both C₂-C₆ alkoxyalkyl and (C₁-C₃)alkoxy(C₁-C₃)alkyl define alkoxyalkyl groups containing from 2 to 6 carbon atoms, the two definitions differ since the former
25 definition allows either the alkoxy or alkyl portion alone to contain 4 or 5 carbon atoms while the latter definition limits either of these groups to 3 carbon atoms.

II. DEFINITIONS

All temperatures are in degrees Centigrade.

TLC refers to thin-layer chromatography.

30 HPLC refers to high pressure liquid chromatography.

Saline refers to an aqueous saturated sodium chloride solution.

Chromatography (column and flash chromatography) refers to purification/separation of compounds expressed as (support, eluent). It is understood that the appropriate fractions are pooled and concentrated to give the desired compound(s).

NMR refers to nuclear (proton) magnetic resonance spectroscopy, chemical shifts are reported in ppm (δ) downfield from tetramethylsilane.

CMR refers to C-13 magnetic resonance spectroscopy, chemical shifts are reported in ppm (δ) downfield from TMS.

5 Pharmaceutically acceptable refers to those properties and/or substances which are acceptable to the patient from a pharmacological/toxicological point of view and to the manufacturing pharmaceutical chemist from a physical/chemical point of view regarding composition, formulation, stability, patient acceptance and bioavailability.

Pharmaceutically acceptable anion salts include salts of the following acids
10 methanesulfonic, hydrochloric, hydrobromic, sulfuric, phosphoric, nitric, benzoic, citric, tartaric, fumaric, maleic, $\text{CH}_3-(\text{CH}_2)_n-\text{COOH}$ where n is 0 thru 4, $\text{HOOC}-(\text{CH}_2)_n-\text{COOH}$ where n is as defined above.

(Z)-2-butenedioate refers to maleate.

When solvent pairs are used, the ratios of solvents used are volume/volume (v/v).

15 When the solubility of a solid in a solvent is used the ratio of the solid to the solvent is weight/volume (wt/v).

Sexual disturbance refers to hypoactive sexual desire disorder, female sexual arousal disorder, male erectile disorder, female orgasmic disorder, and male orgasmic disorder as defined in Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, (DSM-
20 IV), American Psychiatric Association, Washington DC, 1994 as well as the DSM-IV Guidebook, American Psychiatric Press, Inc., Washington DC, 1995. These definitions of the particular disease states are as they were defined in 1994 and 1995. The five "sexual disturbances" are still defined the same today. In the future, the terms for these sexual disturbances may change, but one skilled in the art will know and realize the disease states
25 themselves are the same.

Sexual activity includes sexual intercourse with or without orgasm, ejaculation, masturbation and sexual foreplay.

EXAMPLES

Without further elaboration, it is believed that one skilled in the art can, using the
30 preceding description, practice the present invention to its fullest extent. The following detailed examples describe how to prepare the various compounds and/or perform the various processes of the invention and are to be construed as merely illustrative, and not limitations of the preceding disclosure in any way whatsoever. Those skilled in the art will

promptly recognize appropriate variations from the procedures both as to reactants and as to reaction conditions and techniques.

PREPARATION 1 (R)-Naproxen chloride

R-naproxen (260 g), methylene chloride (3.33 kg) and DMF (8.2 ml) are added to a
5 reactor. Oxalyl chloride (191.8 g) is slowly added to this mixture. After addition of the
oxalyl chloride, the slurry is stirred at 5 to 10° and then slowly warmed to 20-25°. The
resulting mixture is concentrated to remove the methylene chloride, branched octane is
added to the concentrate and the mixture is again concentrated. More branched octane is
added to the concentrate and the mixture is cooled to 0° and stirred to crystallize. The
10 crystal slurry is filtered, the crystal cake is washed with octane and dried at 20-25° to obtain
the title compound.

The filtrate from the first crop is concentrated, branched octane is added and the
mixture is cooled and stirred to obtain a second crop of the title compound. The slurry is
filtered, the crystal cake is washed with branched octane and dried at 20-25°.

15 EXAMPLE 1 1-Benzyl-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (II)

A mixture of 4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (I, *J. Heterocyclic Chem.*, 19,
837-49 (1982), 1.0g, 5.8mmol) in DMF (10ml) is cooled to 0° and treated with potassium *t*-
butoxide in THF (1.98 M, 3.2 ml, 6.3 mmol) maintaining the reaction temperature at 0°.
The resulting mixture is stirred at 0° for 10 minutes. Benzyl bromide (0.73 ml, 6.1mmol) is
20 then added while maintaining the reaction temperature at 0°. After 1 hr, the mixture is
partitioned with methyl *t*-butyl ether (MTBE) from water followed by several water washes.
The MTBE phase is concentrated under reduced pressure. The concentrate is cooled to 0°,
filtered and washed two times with 0° MTBE. The product is dried at 50° under reduced
pressure with a nitrogen purge to give the title compound, CMR (CDCl₃, 100 MHz) L
25 153.78, 136.44, 128.69, 127.67, 127.60, 126.73, 125.86, 122.90, 122.78, 121.28, 116.92,
116.17, 108.36, 44.95 and 42.37.

EXAMPLE 2 (5R*,6R*)-1-benzyl-5-bromo-6-hydroxy-5,6-dihydro-4H- imidazo[4,5,1-ij]quinolin-2(1H)-one (III)

1-Benzyl-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (II, XAMPLE 1, 240 g),
30 acetonitrile (1.086 kg), water (227 ml) and fluoboric acid (48.5%, 13.4 g) are mixed and
cooled to 0 to 5°. Dibromantin (163.5 g) is slurried into acetonitrile and is added to the
reaction mixture. The reaction is carried out for about 3 hr at 0 to 5°. After the reaction is
complete, methyl *t*-butyl ether is added over about 45 minutes keeping the reaction

temperature in the pot below 10°. The slurry is cooled to -10 to -15°, stirred for an hour and then filtered. The product is washed with precooled methyl *t*-butyl ether, dried with 40° nitrogen to give the title compound, CMR (CDCl₃) δ 156.0, 137.8, 130.5, 129.6, 129.3, 129.1, 126.6, 123.6, 122.5, 119.6, 110.4, 69.9, 49.6, 47.7, 46.9 and 43.8.

- 5 **EXAMPLE 3** (5S,6S)-1-Benzyl-5-bromo-2-oxo-1,2,5,6-tetrahydro-4H-imidazo[4,5,1-*ij*]quinolin-6-yl (2R)-2-(6-methoxy-2-naphthyl)propanoate (IVA) and (5R,6R)-1-benzyl-5-bromo-2-oxo-1,2,5,6-tetrahydro-4H-imidazo[4,5,1-*ij*]quinolin-6-yl (2R)-2-(6-methoxy-2-naphthyl)propanoate (IVB)
- 10 (5R,6R)-1-Benzyl-5-bromo-6-hydroxy-5,6-dihydro-4H-imidazo[4,5,1-*ij*]quinolin-2(1H)-one (III, EXAMPLE 2, 143 g), methylene chloride (3,136 g), N-methyl morpholine (100.2 g) and 4-dimethylaminopyridine (497 mg) are added to the reactor and the mixture is cooled to 0 to 5°. (R)-Naproxen chloride (PREPARATION 1, 118.5 g) dissolved in methylene chloride (694 ml) is added to the reactor over about 1 hr and the mixture is
- 15 stirred at 0 to 5° to complete the reaction. If necessary, additional naproxen chloride is added to complete the reaction. Potassium carbonate solution diluted with water is added to the mixture. The aqueous phase is extracted with methylene chloride and the combined methylene phase is washed with water. The washed mixture is concentrated by vacuum distillation and solvent exchange with ethyl acetate is performed. The concentrate is cooled
- 20 to -10° and stirred. The crystal slurry is filtered and the crystal cake is washed with precooled methyl *t*-butyl ether and dried at 50° to give the title compound in solid form, (5S,6S)-1-benzyl-5-bromo-2-oxo-1,2,5,6-tetrahydro-4H-imidazo[4,5,1-*ij*]quinolin-6-yl (2R)-2-(6-methoxy-2-naphthyl)propanoate (IVA), CMR (CDCl₃) δ 173.2, 157.8, 153.4, 136.1, 134.6, 133.7, 129.2, 128.8, 127.8, 127.8, 127.6, 127.2, 125.9, 125.9, 125.6, 121.5,
- 25 121.4, 119.1, 113.2, 109.0, 105., 105.6, 69.2, 55.3, 45.4, 45.2, 42.5, 41.7 and 18.3.

The undesired isomer, (5R,6R)-1-benzyl-5-bromo-2-oxo-1,2,5,6-tetrahydro-4H-imidazo[4,5,1-*ij*]quinolin-6-yl (2R)-2-(6-methoxy-2-naphthyl)propanoate (IVB) is in the filtrate and can be recovered by means well known to those skilled in the art, (5R,6R)-1-benzyl-5-hydroxy-6-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-*ij*]quinolin-2(1H)-one,

30 CMR 173.2, 157.9, 153.4, 136.1, 135.0, 133.8, 129.2, 128.9, 128.8, 127.8, 127.6, 127.4, 125.8, 125.8, 125.7, 121.6, 121.5, 119.3, 113.1, 109.1, 105.7, 68.7, 55.3, 45.3, 45.2, 42.2, 41.3 and 18.1 δ.

EXAMPLE 4 (5R,6R)-1-benzyl-5-hydroxy-6-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (V)

(5S,6S)-1-Benzyl-5-bromo-2-oxo-1,2,5,6-tetrahydro-4H-imidazo[4,5,1-ij]quinolin-6-yl (2R)-2-(6-methoxy-2-naphthyl)propanoate (IVA, EXAMPLE 3, 110 g) is slurried in acetonitrile (1,297 g). After adding aqueous methylamine (40 wt %, 327 g) the reaction is carried out for about 12 hr at about 30°. After the reaction is complete, the mixture is concentrated and ethyl acetate is added. Dilute hydrochloric acid is added to make the water-soluble salt of the title compound. The byproduct (R-naproxen acetamide impurity) is insoluble in water and stays in the ethyl acetate phase. Further extractions and washes are carried out for better separation of the (naproxen acetamide) impurity with minimum loss of the desired product. Then a sodium hydroxide solution is added to the aqueous phase and the hydrochloride salt of the title compound is converted to the free base. The free base is less soluble in water and is extracted into ethyl acetate. The product mixture is concentrated and solvent exchanged with ethyl acetate to remove water. Crystallization is performed by adding branched chain octane and cooling the mixture. The resulting slurry is filtered, washed and dried at 50° to give the title compound, CMR (CDCl₃) δ 153.7, 136.3, 128.7, 127.8, 127.7, 125.7, 121.3, 119.9, 118.6, 107.5, 66.2, 60.1, 45.1, 42.6 and 34.0.

EXAMPLE 5 (7aS,8aR)-4-benzyl-8-methyl-7,7a,8,8a-tetrahydroazireno[2,3-c]imidazo[4,5,1-ij]quinolin-5(4H)-one (VI)

(5R,6R)-1-benzyl-5-hydroxy-6-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (V, EXAMPLE 4, 70 g) and THF (1,389 g) is concentrated to remove any moisture with distillate as a precaution due to reactivity of *n*-butyl lithium towards water. The mixture is cooled to about -10° and *n*-butyllithium is added to make the lithium salt of the starting material with formation of *n*-butane byproduct in an exothermic reaction. Benzene sulfonyl chloride is added slowly to make benzene sulfonate in an exothermic reaction. The reaction mixture is warmed to 20-25° to complete the reaction. Aqueous potassium carbonate solution is added to scavenge the benzene sulfonic acid and the mixture is stirred to allow crystallization. Water is added to complete crystallization, the slurry is stirred, cooled and filtered. The crystal cake is washed with water followed by branched chain octane and dried at 40 to 50° to give the title compound, CMR (CDCl₃) δ 154.1, 136.3, 128.6, 127.9, 127.6, 124.3, 120.7, 119.7, 107.4, 46.7, 44.9, 40.7, 38.1 and 37.6.

EXAMPLE 6 (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (VII)

A mixture of (7aS,8aR)-4-benzyl-8-methyl-7,7a,8,8a-tetrahydroazireno[2,3-c]imidazo[4,5,1-ij]quinolin-5(4H)-one (VI, EXAMPLE 5, 40 g) *t*-amyl alcohol (42.4 g) and
5 anhydrous ammonia (1,200 g) is treated with lithium at -33°. After the lithium addition is complete, the reaction mixture changes from a yellow slurry to a dark blue mixture. This dark blue mixture is stirred for 30-60 minutes and then quenched with the addition of water. The cooling is removed from the condenser and the ammonia is allowed to evaporate. The residue is dissolved in methanol. This mixture is then concentrated to dryness to give the
10 title compound, which is carried on directly to the next step without isolation.

EXAMPLE 7 (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (Z)-2-butenedioate (1:1) (VIII)

(5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (VII, EXAMPLE 6, 28.0 g) is dissolved in water and the pH is adjusted to 10 with the addition of
15 hydrochloric acid. The mixture is applied in portions to an XAD-16 resin column which is eluted first with water and then with ethanol. The inorganic salts are eluted from the column first with the desired product eluted with the ethanol. The ethanol eluate from the column is treated with maleic acid and the water level is lowered through azeotropic distillation of the ethanol. The precipitated product is isolated by filtration, rinsed with
20 ethyl acetate and dried to give the title compound, CMR (DMSO-*d*₆) δ 167.6, 153.9, 136.4, 127.1, 121.5, 119.6, 114.1, 107.5, 51.9, 31.3 and 26.5.

EXAMPLE 8 (5R)-5-(Methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinoline-2(1H)-thione

A mixture of (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one
25 (VII, EXAMPLE 6, 15.0 g, 73.8 mmol) and tetraphosphorus decasulfide (36.1 g, 81.2 mmol) in pyridine (300 mL) is heated in a 125° oil bath under nitrogen. The reaction is stirred for 5 hr. The mixture is cooled to 20-25° and the pyridine is removed under reduced pressure. Sodium hydroxide (2.2 N, 200 mL) is added. Sodium hydroxide (1 N) is then added. The mixture is saturated with sodium chloride and extracted with methylene
30 chloride (2.5 L, in portions). The organic phase is absorbed onto silica gel (40 g) and purified via column chromatography (silica gel; 225 g; methanol/methylene chloride, 3.5-5.0/96.5-95) to give a solid. Recrystallization of this material from methanol/ethyl acetate/hexanes give the title compound, mp = 210-213°; IR (drift) 2940, 2907, 2884,

1483, 1458, 1391, 1366, 1354, 1254, 1239, 1229, 895, 762, 734, 630 cm^{-1} ; NMR (300 MHz, CDCl_3) 7.12, 7.03, 7.00, 4.30, 3.96, 3.30-3.50, 3.15, 2.88 and 2.57 δ ; MS (EI, m/z) 219 (M^+), 190, 189, 187, 186, 164, 163, 155, 145; HRMS (FAB) calculated for $\text{C}_{11}\text{H}_{13}\text{N}_3\text{S}$ (MH^+) = 220.0908, found 220.0904.

5 **EXAMPLE 9** (5R)-5-(Methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinoline-2(1H)-thione malate

A mixture of maleic acid (0.317 g, 2.36 mmol) in a minimal amount of methanol (~ 1 mL) is added to a mixture of (5R)-5-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinoline-2(1H)-thione (EXAMPLE 8, 0.493 g, 2.25 mmol) in methylene chloride. The resulting
10 solid is collected by filtration to give the title compound, mp = 195-196°; $[\alpha]_{\text{D}}^{25} = -60^\circ$ (c 0.93, methanol); IR (drift) 3140, 3112, 3060, 2969, 1627, 1619, 1568, 1481, 1455, 1398, 1389, 1361, 1220, 868 and 747 cm^{-1} ; NMR (300 MHz, CD_3OD) 7.20-7.30, 7.10-7.20, 6.26, 4.49, 4.31, 4.05-4.20, 3.47, 3.28 and 2.83 δ ; CMR (100 MHz, $\text{DMSO}-d_6 + \text{CD}_3\text{OD}$) 170.4, 169.4, 136.6, 131.1, 130.9, 125.1, 122.1, 116.2, 109.6, 53.9, 43.1, 31.9 and 27.2 δ ; MS
15 (ESI, m/z) 220.1 (MH^+).

EXAMPLE A Hypoactive Sexual Desire Disorder In A Female Patient

A 44 kg, 24 year old female patient is referred by her psychologist for evaluation for and medical treatment of hypoactive sexual desire disorder. The patient recently began to co-habitate with a 28 year old man. She consulted a clinical psychologist because of
20 distress over frequent arguments with her partner. These arguments concern her frustration that her partner is unwilling to marry her and the partner's accusations that she is sexually unresponsive and "boring in bed." The patient reports that although she desires to continue the relationship, she has little or no interest in sexual activity with this partner. The partner initiates all sexual activity in the relationship and she complies so that the relationship will
25 continue. She states that her partner desires sexual intercourse about once per day. Occasionally, the couple's frequency of sexual intercourse is as much as twice per day. She reports that she does not have sexual fantasies since the present relationship began. The patient had a previous relationship with a man that lasted 3 years and involved sexual activity that she found sexually exciting and satisfying. The patient has recently had a
30 physical examination by her family physician, the findings of which were unremarkable. She currently takes no prescription drugs other than low dose birth control pills. She does not routinely use over-the-counter medications or alcohol and denies taking drugs of abuse. The patient is diagnosed as having Hypoactive Sexual Desire Disorder of acquired onset due to situational factor (Diagnostic and Statistical Manual of Mental Disorders, Fourth

Edition, 1994). (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (Z)-2-butenedioate (1:1) is prescribed at 1 mg/dose to be taken 0.5 hr before the patient intends to engage in sexual activity. The patient is instructed that she may use the drug on a daily basis, if she and her partner desire sexual activity at this frequency. She is also
5 instructed that on occasions she may use the drug twice in a single day if this is the intended frequency of intercourse. The patient is examined after one month. She reports that she uses the drug once, and occasionally twice, per day. She reports that she has sexual fantasies on 3-4 occasions per week, occasionally initiates sexual activity with her partner, and enjoys an improved sexual relationship with her partner. She also reports that she and
10 her partner argue less frequently and that her partner no longer accuses her of being disinterested in sexual relations. The patient is instructed to continue using the drug.

EXAMPLE B Female Sexual Arousal Disorder In A Female Patient

A 64 kg, 22 year old female patient is complaining of vaginal dryness and mild discomfort during sexual intercourse. The patient recently began a relationship with a 35
15 year old man. Her partner complained of her vaginal dryness and poor vaginal expansion during sexual intercourse. Her partner has made several comments about this during the previous two months and the patient is concerned that her partner is dissatisfied with her lack of sexual responsiveness and may end the relationship. The patient reports that she desires to continue the relationship. The patient has had several previous sexual
20 relationships. She reports that vaginal dryness and some discomfort with vaginal intercourse has occurred during her past sexual relationships. On questioning, she indicates that she has little or no sense of sexual arousal during intercourse. The patient indicates that she has sexual fantasies several times per week. She finds her current partner attractive and sexual desirable. She indicates that she masturbates approximately once per week.
25 During these episodes she usually has an orgasm. Lubrication is not a problem when she masturbates. The patient has recently had a physical examination by her family physician, the findings of which were unremarkable. She currently takes no prescription drugs other than low dose birth control pills. She does not routinely use over-the-counter medications or alcohol and denies taking drugs of abuse. The patient is diagnosed as having Female
30 Sexual Arousal Disorder that is situational due to psychological factors (Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, 1994). (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (Z)-2-butenedioate (1:1) is prescribed at 3 mg/dose to be taken 0.5-1 hr before the patient intends to engage in sexual activity. The patient is instructed that she may use the drug on a daily basis, if this is the intended

frequency of sexual activity. The patient is examined after one month. She reports that she uses the drug prior to intercourse. She reports that vaginal dryness is no longer a problem and that she no longer has pain during initiation of vaginal penetration by her partner. She enjoys an improved sexual relationship with her partner. The patient is instructed to

5 continue using the drug.

EXAMPLE C Male Erectile Disorder

A 75 kg, 56 year old male patient has been married for 20 years. He presents complaining that for the past six months he has had difficulty maintaining an erection until completion of sexual intercourse. He develops an adequate erection at the beginning of
10 sexual activity but detumescence occurs when he attempts penetration. His wife is his only current sexual partner. He reports that his wife is patient and understanding of his problem but that he is embarrassed, feels that he is a failure, and is growing progressively more angry and sad about his inability to maintain an erection. He masturbates approximately twice each month. During masturbation, he is usually able to maintain an erection until he
15 ejaculates. The patient has recently had a physical examination by his family physician. The only remarkable findings were a supine diastolic blood pressure 94 mm mercury and a total plasma cholesterol level of 230 mg/dl. He was instructed to follow the American Heart Association dietary recommendations and to reduce his body weight by 5 kg. He currently takes no prescription drugs. He does not routinely use over-the-counter
20 medications or alcohol and denies taking drugs of abuse. The patient is diagnosed as having Male Erectile Disorder that is acquired and situational. The etiology is most likely due to psychological factors, although combined factors, e.g., mild hypertension or atherosclerosis, may contribute (Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, 1994). (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one
25 (Z)-2-butenedioate (1:1) is prescribed at 3 mg/dose to be taken 0.5 hr before the patient intends to engage in sexual activity. The patient is instructed that he may use the drug on a daily basis, if this is the intended frequency of sexual activity. The patient is examined after one month. He reports that he uses the drug prior to intercourse. He reports that he maintains an erection for a satisfactory period after vaginal penetration in almost every
30 attempt at intercourse. He usually experiences intra-vaginal ejaculation. The patient and his wife consider the quality of their sexual relationship to be markedly improved. The patient is instructed to continue using the drug.

EXAMPLE D Female Orgasmic Disorder

A 50 kg, 22 year old female patient who married three months before presenting for treatment. She complains that she does not have an orgasm during sexual activity with her husband even though she enjoys the experience and becomes sexually excited during intercourse. She was not sexually active prior to marriage. She reports that her husband
5 had multiple premarital partners and that he reports all his past partners had orgasms. The patient is upset by her inability to have orgasms and feels she is a poor wife. On questioning she reports that she is "too fat" although her weight is appropriate for her height. She does not masturbate. The patient has recently had a physical examination by her family physician, the findings of which were unremarkable. She currently takes no
10 prescription drugs other than low dose birth control pills. She does not routinely use over-the-counter medications or alcohol and denies taking drugs of abuse. The patient is diagnosed as having Female Orgasmic Disorder that is situational due to psychological factors (Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, 1994).
(5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (Z)-2-butenedioate (1:1)
15 is prescribed at 1 mg/dose to be taken about 0.5 hr before the patient intends to engage in sexual activity. The patient is instructed that she may use the drug on a daily basis, if this is the intended frequency of sexual activity. The patient is examined after two months. She reports that she uses the drug prior to intercourse. She reports that she has an orgasm during sexual intercourse on most occasions. She enjoys an improved sexual relationship
20 with her partner. It is recommended that the patient continue to use the drug for an additional six months, then stop taking the drug for a period of two weeks to determine whether she might now be able to have orgasms in the absence of drug therapy.

EXAMPLE E Male Orgasmic Disorder

A 80 kg, 40 year old man has been married for 15 years. He presents complaining
25 that he has begun to have difficulty achieving an orgasm during vaginal intercourse with his wife. On most occasions he is unable to ejaculate during sexual intercourse. Infrequently, he is able to ejaculate during sexual, but only after very prolonged vaginal intercourse. He indicates that this has been occurring for at least three months and that he is frustrated and angry about the situation. His further reports that his wife finds the prolonged intercourse
30 that he requires to ejaculate uncomfortable and has begun avoiding sexual relations. He reports that he ejaculates normally when he masturbates. The patient has recently had a physical examination by his family physician, the findings of which were unremarkable. He currently takes no prescription drugs. He does not routinely use over-the-counter medications or alcohol and denies taking drugs of abuse. The patient is diagnosed as

having Male Orgasmic Disorder that is acquired and situational due to psychological factors (Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, 1994). (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (Z)-2-butenedioate (1:1) is prescribed at 3 mg/dose to be taken 0.5-1 hr before the patient intends to engage in sexual activity. The patient is instructed that he may use the drug on a daily basis, if this is the intended frequency of sexual activity. The patient is examined after one month. At this visit he reports that he now ejaculates during vaginal intercourse on all occasions. The patient and his wife consider the quality of their sexual relationship to be markedly improved. The patient is instructed to continue using the drug.

10 **EXAMPLE F Mating of Animals of Commercial Value**

 An adult male Giant Panda on temporary loan from a foreign zoo is desired to have breed to an adult female Giant Panda which is normally resident at the zoo. Because of the commercial value of a potential off-spring it is desirable that the animals mate during the time the adult male is on loan. Both animals are sexually inexperienced and do not engage in intercourse during a two week period. As an alternative to anesthetizing the animals for a facilitated transfer of semen between the animals, the male and female Giant Pandas have (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (Z)-2-butenedioate (1:1) added to their diet on one occasion. The concentration of drug in the diet is calculated to deliver 0.05 mg of free base equivalents of (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (Z)-2-butenedioate (1:1) per kilogram of body weight. One hour after ingesting the diet containing drug, the animals engage in sexual intercourse. A pregnancy and subsequent birth of an off-spring results from this intercourse.

EXAMPLE G Reduced Sexual Function In A Male With Idiopathic Epilepsy

 A 40 year old, 75 kg male patient has idiopathic epilepsy. He first presented with primary generalized seizures at age 15 years and was diagnosed with idiopathic epilepsy. He has been treated with carbamazepine since age 20. Currently he receives 500 mg/day of the drug. He has had two tonic-clonic seizures during the preceding year. The patient complains of diminished sexual interest in his wife of 8 years. On those occasions when he attempts sexual intercourse, he is able to maintain an erection sufficient for intercourse one-third of the time. He indicates that this problem has had a gradual onset over the previous 5 years. He reports that he rarely has sexual fantasies and rarely masturbates. The patient has recently had a physical examination by his family physician, the findings of which were unremarkable. He currently takes no prescription drugs other than carbamazepine. He does not routinely use over-the-counter medications or alcohol and denies taking drugs of abuse.

A penile injection test with Caverject™ Injection results in a normal erection and duplex Doppler ultrasonography indicates an intact vascular system. Plasma testosterone and sex hormone binding globulin assays are performed. The plasma testosterone level is 25 nM, which is considered normal. The plasma sex hormone binding globulin level is 55 nM, which is considered elevated compared to the normal range. The patient is diagnosed with hypogonadism, secondary to chronic carbamazepine treatment (Epilepsia 36: 366-370, 1995). (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (Z)-2-butenedioate (1:1) is prescribed at 3 mg/dose to be taken 0.5 hr before the patient intends to engage in sexual activity. The patient is instructed that he may use the drug on a daily basis, if this is the intended frequency of sexual activity. The patient is examined after one month. He reports that he take the drug 2-3 times per week. After taking the drug, he feels more interest in sex and usually initiates sexual activity with his wife. He reports that enjoys a normal sexual relationship. The patient is instructed to continue with the treatment.

EXAMPLE H Reduced Sexual Function In A Male Patient With Erectile Dysfunction And Hypoactive Sexual Desire Disorder

A 55 year old, 70 kg, male patient is apparently in good health. He presents complaining of a loss of sexual desire and frequent erectile dysfunction since taking an executive position in his company, six weeks previously. He reports that he works long hours and feels a high degree of pressure associated with his job. He is frustrated and angry that when he takes time off from work he is unable to have an intimate relationship with his wife. He reports that although he is periodically interested in initiating sexual relations, on most occasions he is unable to maintain a high level of interest sufficiently long to initiate or to adequately respond to his wife's initiating sexual activity. He further indicates that for the past six months, he has had difficulty maintaining an erection until completion of sexual intercourse. He develops a semi-rigid erection at the beginning of sexual activity but detumescence occurs when he attempts penetration. He reports that he has few erotic fantasies. He has not masturbated for the past several months. He attributes this to a lack of motivation and interest. The patient has recently had a physical examination by his family physician, the findings of which were unremarkable. He does not routinely use over-the-counter medications or alcohol and denies taking drugs of abuse. (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (Z)-2-butenedioate (1:1), 3 mg/dose, and sildenafil (VIAGRA Tablets), 50 mg/dose, are both prescribed. Each is to be taken orally about 0.5 hr before the patient intends to engage in sexual activity. The patient is

instructed that he may use the drug combination on a daily basis, if this is the intended frequency of sexual activity. The patient is examined after one month. He reports that he takes the drug combination 2-3 times per week. After taking the drug combination he feels motivated to initiate sexual activity, readily has sexual fantasies that are sustained during sexual intercourse, and has no difficulty maintaining an erection to completion of sexual intercourse. He reports that he enjoys an improved sexual relationship with his partner. The patient is advised to continue taking the (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (Z)-2-butenedioate (1:1) for another six months as he adapts to the situational stress associated with his new job. After this time he is advised to stop taking the drug for a period of two weeks to determine whether his sexual motivation and desire have returned to the level he enjoyed prior to the change in jobs.

EXAMPLE I Hypoactive Sexual Desire Disorder In A Female Patient

Following the general method of treatment of EXAMPLE A and making non-critical variations but using (5R)-5-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinoline-2(1H)-thione in place of (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (Z)-2-butenedioate (1:1) the same positive treatment effect is obtained.

EXAMPLE J Female Sexual Arousal Disorder In A Female Patient

Following the general method of treatment of EXAMPLE B and making non-critical variations but using (5R)-5-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinoline-2(1H)-thione in place of (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (Z)-2-butenedioate (1:1) the same positive treatment effect is obtained.

EXAMPLE K Male Erectile Disorder

Following the general method of treatment of EXAMPLE C and making non-critical variations but using (5R)-5-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinoline-2(1H)-thione in place of (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (Z)-2-butenedioate (1:1) the same positive treatment effect is obtained.

EXAMPLE L Female Orgasmic Disorder

Following the general method of treatment of EXAMPLE D and making non-critical variations but using (5R)-5-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinoline-2(1H)-thione in place of (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (Z)-2-butenedioate (1:1) the same positive treatment effect is obtained.

EXAMPLE M Male Orgasmic Disorder

Following the general method of treatment of EXAMPLE E and making non-critical variations but using (5R)-5-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinoline-2(1H)-thione in place of (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (Z)-2-butenedioate (1:1) the same positive treatment effect is obtained.

5 **EXAMPLE N Mating of Animals of Commercial Value**

Following the general method of treatment of EXAMPLE F and making non-critical variations but using (5R)-5-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinoline-2(1H)-thione in place of (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (Z)-2-butenedioate (1:1) the same positive treatment effect is obtained.

10 **EXAMPLE O Reduced Sexual Function In A Male With Idiopathic Epilepsy**

Following the general method of treatment of EXAMPLE G and making non-critical variations but using (5R)-5-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinoline-2(1H)-thione in place of (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (Z)-2-butenedioate (1:1) the same positive treatment effect is
15 obtained.

**EXAMPLE P Reduced Sexual Function In A Male Patient With Erectile
 Dysfunction And Hypoactive Sexual Desire Disorder**

Following the general method of treatment of EXAMPLE H and making non-critical variations but using (5R)-5-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-
20 ij]quinoline-2(1H)-thione in place of (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (Z)-2-butenedioate (1:1) the same positive treatment effect is
obtained.




CHART A






CHART A - continued


CHART A - continued



5

CLAIMS

1. A method of treating sexual disturbances in a human who is in need of such treatment which comprises administering a sexually therapeutically effective amount of a compound of the formula (A)



5

where

R₁, R₂ and R₃ are the same or different and are:

-H,

C₁-C₆ alkyl,

10

C₃-C₅ alkenyl,

C₃-C₅ alkynyl,

C₃-C₅ cycloalkyl,

C₄-C₁₀ cycloalkyl,

phenyl substituted C₁-C₆ alkyl,

15

-NR₁R₂ where R₁ and R₂ are cyclized with the attached nitrogen atom to produce pyrrolidyl, piperidyl, morpholinyl, 4-methyl piperazinyl or imidazolyl;

X is:

-H,

C₁-C₆ alkyl,

20

-F, -Cl, -Br, -I,

-OH,

C₁-C₆ alkoxy,

cyano,

carboxamide,

25

carboxyl,

(C₁-C₆ alkoxy)carbonyl,

A is:

CH,

CH₂,

CH-(halogen) where halogen is -F, -Cl, -Br, -I,

CHCH₃,

C=O,

C=S,

5 C-SCH₃,

C=NH,

C-NH₂,

C-NHCH₃,

C-NHCOOCH₃,

10 C-NHCN,

SO₂,

N;

B is:

CH₂,

15 CH,

CH-(halogen) where halogen is as defined above,

C=O,

N,

NH,

20 N-CH₃,

D is:

CH,

CH₂,

CH-(halogen) where halogen is as defined above,

25 C=O,

O,

N,

NH,

N-CH₃;

30 and n is 0 or 1, and where \sim is a single or double bond, with the provisos:

(1) that when n is 0, and

A is CH₂, CH-(halogen) where halogen is as defined above, CHCH₃, C=O, C=S, C=NH, SO₂;

then D is CH₂, CH-(halogen) where halogen is as defined above, C=O, O, NH, N-CH₃;

(2) that when n is 0, and

A is CH, C-SCH₃, C-NH₂, C-NHCH₃, C-NHCOOCH₃, C-NHCN, N; then

5 D is CH, N;

(3) that when n is 1, and

A is CH₂, CH-(halogen) where halogen is as defined above, CHCH₃, C=O, C=S, C=NH, SO₂; and

10 B is CH₂, CH-(halogen) where halogen is as defined above, C=O, NH, N-CH₃; then

D is CH₂, C=O, O, NH, N-CH₃;

(4) that when n is 1, and

A is CH, C-SCH₃, C-NH₂, C-NHCH₃, C-NHCOOCH₃, C-NHCN, N; and

B is CH, N; then

15 D is CH₂, C=O, O, NH, N-CH₃;

(5) that when n is 1, and

A is CH₂, CHCH₃, C=O, C=S, C=NH, SO₂, and

B is CH, N; then

D is CH, N; and pharmaceutically acceptable salts thereof to the human.

20

2. A method of treating sexual disturbances according to claim 1 where the mammal is a male.

25 3. A method of treating sexual disturbances according to claim 1 where the mammal is a female.

4. A method of treating sexual disturbances according to claim 1 where the sexual disturbance is selected from the group consisting of hypoactive sexual desire disorder, female sexual arousal disorder, male erectile disorder, female orgasmic disorder, and male
30 orgasmic disorder.

5. A method of treating sexual disturbances according to claim 4 where the sexual disturbance is hypoactive sexual desire disorder.

6. A method of treating sexual disturbances according to claim 4 where the sexual disturbance is female sexual arousal disorder.
7. A method of treating sexual disturbances according to claim 4 where the sexual
5 disturbance is male erectile disorder.
8. A method of treating sexual disturbances according to claim 4 where the sexual disturbance is female orgasmic disorder.
- 10 9. A method of treating sexual disturbances according to claim 4 where the sexual disturbance is male orgasmic disorder.
10. A method of treating sexual disturbances according to claim 1 where the compound of formula (A) is administered orally, intra-nasally, buccally, intra-pulmonary, parenterally
15 and rectally.
11. A method of treating sexual disturbances according to claim 10 where the compound of formula (A) is administered orally, intra-nasally, buccally and intra-pulmonary.
- 20 12. A method of treating sexual disturbances according to claim 10 where the compound of formula (A) is administered orally.
13. A method of treating sexual disturbances according to claim 1 where the sexually therapeutically effective amount is from about 0.2 thru about 8 mg/person/dose.
25
14. A method of treating sexual disturbances according to claim 13 where the sexually therapeutically effective amount is from about 0.5 thru about 5 mg/person/dose.
15. A method of treating sexual disturbances according to claim 14 where the sexually
30 therapeutically effective amount is from about 1 thru about 3 mg/person/dose.
16. A method of treating sexual disturbances according to claim 1 where the compound of formula (A) is (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one.

17. A method of treating sexual disturbances according to claim 16 where the compound of formula (A) is (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (Z)-2-butenedioate (1:1).

5 18. A method of treating sexual disturbances according to claim 1 where the pharmaceutically acceptable salt is selected from the group consisting of salts of the following acids methanesulfonic, hydrochloric, hydrobromic, sulfuric, phosphoric, nitric, benzoic, citric, tartaric, fumaric, maleic, $\text{CH}_3-(\text{CH}_2)_n-\text{COOH}$ where n is 0 thru 4, $\text{HOOC}-(\text{CH}_2)_n-\text{COOH}$ where n is as defined above.

10

19. A method of treating sexual disturbances according to claim 1 where the compound of formula (A) is administered from about 10 minutes to about 8 hr prior to sexual activity.

20. A method of treating sexual disturbances according to claim 19 where the compound of
15 formula (A) is administered from about 0.5 hr to about 1 hr prior to sexual activity.

21. A method of treating sexual disturbances according to claim 20 where the compound of formula (A) is administered about 0.5 hr prior to sexual activity.

20 22. A method of treating sexual disturbances according to claim 1 where the mammal does not have Parkinson's disease.

23. A method of treating sexual disturbances according to claim 1 where the mammal does not experience postural hypotension.

25

24. A method of treating sexual disturbances according to claim 1 where the compound of formula (A) is used in combination with a sexually effective amount of one or more vascular smooth muscle relaxation agents where both the compound of formula (A) is administered within 8 hours prior to sexual activity and the vascular smooth muscle
30 relaxation agent is administered to the human within a sexually effective time period prior to sexual activity.

25. A method of treating sexual disturbances according to claim 24 where the vascular smooth muscle relaxation agents is selected from the group consisting of phosphodiesterase


type 5 inhibitors, phosphodiesterase type 3 inhibitors, non-selective phosphodiesterase inhibitors, nitric oxide donor drugs, alpha type 1 adrenergic receptor antagonists, alpha type 2 adrenergic receptor antagonists, prostaglandin E1 receptor agonists (PGE1) and vasoactive intestinal polypeptide (VIP) agents.

5

26. A method of treating sexual disturbances according to claim 25 where the vascular smooth muscle relaxation agents is selected from the group consisting of sildenafil, ICOS-351, milrinone, papaverine, linsidomine, phentolamine, yohimbine, prostaglandin E1 (PGE1) and VIP.

10

27. A method of inducing mating a non-human mammal which comprises administering a sexually mating amount of a compound of the formula (A)



where

15 R_1 , R_2 and R_3 are the same or different and are:

-H,

C_1 - C_6 alkyl,

C_3 - C_5 alkenyl,

C_3 - C_5 alkynyl,

20 C_3 - C_5 cycloalkyl,

C_4 - C_{10} cycloalkyl,

phenyl substituted C_1 - C_6 alkyl,

- NR_1R_2 where R_1 and R_2 are cyclized with the attached nitrogen atom to produce pyrrolidyl, piperidinyl, morphoninyl, 4-methyl piperazinyl or imidazolyl;

25 X is:

-H,

C_1 - C_6 alkyl,

-F, -Cl, -Br, -I,

-OH,

C₁-C₆ alkoxy,
cyano,
carboxamide,
carboxyl,
5 (C₁-C₆ alkoxy)carbonyl,

A is:

CH,
CH₂,
CH-(halogen) where halogen is -F, -Cl, -Br, -I,
10 CHCH₃,
C=O,
C=S,
C-SCH₃,
C=NH,
15 C-NH₂,
C-NHCH₃,
C-NHCOOCH₃,
C-NHCN,
SO₂,
20 N;

B is:

CH₂,
CH,
CH-(halogen) where halogen is as defined above,
25 C=O,
N,
NH,
N-CH₃,

D is:

30 CH,
CH₂,
CH-(halogen) where halogen is as defined above,
C=O,
O,

N,
NH,
N-CH₃;

and n is 0 or 1, and where --- is a single or double bond, with the provisos:

5 (1) that when n is 0, and

A is CH₂, CH-(halogen) where halogen is as defined above, CHCH₃, C=O, C=S, C=NH, SO₂;

then D is CH₂, CH-(halogen) where halogen is as defined above, C=O, O, NH, N-CH₃;

10 (2) that when n is 0, and

A is CH, C-SCH₃, C-NH₂, C-NHCH₃, C-NHCOOCH₃, C-NHCN, N; then D is CH, N;

(3) that when n is 1, and

A is CH₂, CH-(halogen) where halogen is as defined above, CHCH₃, C=O, C=S, C=NH, SO₂; and

B is CH₂, CH-(halogen) where halogen is as defined above, C=O, NH, N-CH₃; then

D is CH₂, C=O, O, NH, N-CH₃;

(4) that when n is 1, and

20 A is CH, C-SCH₃, C-NH₂, C-NHCH₃, C-NHCOOCH₃, C-NHCN, N; and B is CH, N; then

D is CH₂, C=O, O, NH, N-CH₃;

(5) that when n is 1, and

A is CH₂, CHCH₃, C=O, C=S, C=NH, SO₂, and

25 B is CH, N; then


D is CH, N; and pharmaceutically acceptable salts thereof.

28. A method of inducing mating according to claim 27 where the non-human mammal is selected from the group consisting of horses, cattle, swine, sheep, transgenic mice, panda
30 bears, elephants, zebras, lions, tigers, monkeys, apes, dogs and cats.

29. A method of inducing mating according to claim 27 where the non-human mammal is a male.

30. A method of inducing mating according to claim 27 where the non-human mammal is a female.
31. A method of inducing mating according to claim 27 where the compound of formula
5 (A) is administered orally, parenterally and rectally.
32. A method of inducing mating according to claim 31 where the compound of formula (A) is administered orally.
- 10 33. A method of inducing mating according to claim 27 where the sexually mating amount is from about 0.003 thru about 0.2 mg/kg/dose.
34. A method of inducing mating according to claim 33 where the sexually mating amount is from about 0.01 thru about 0.125 mg/kg/dose.
- 15 35. A method of inducing mating according to claim 27 where the sexually mating amount is from about 0.025 thru about 0.075 mg/kg/dose.
36. A method of inducing mating according to claim 27 where the compound of formula
20 (A) is (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one.
37. A method of inducing mating according to claim 36 where the compound of formula (A) is (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (Z)-2-butenedioate (1:1).
- 25 38. A method of inducing mating according to claim 27 where the pharmaceutically acceptable salt is selected from the group consisting of salts of the following acids methanesulfonic, hydrochloric, hydrobromic, sulfuric, phosphoric, nitric, benzoic, citric, tartaric, fumaric, maleic, $\text{CH}_3-(\text{CH}_2)_n-\text{COOH}$ where n is 0 thru 4, $\text{HOOC}-(\text{CH}_2)_n-\text{COOH}$
30 where n is as defined above.
39. A method of inducing mating according to claim 27 where the compound of formula (A) is administered from about 10 minutes to about 8 hr prior to mating.

40. A method of inducing mating according to claim 39 where the compound of formula (A) is administered from about 10 minutes to about 1 hr prior to mating.
41. A method of inducing mating according to claim 40 where the compound of formula (A) is administered from about 10 minutes to about 0.5 hr prior to mating.
42. A method of inducing mating according to claim 27 where the compound of formula (A) is used in combination with a sexually effective amount of one or more vascular smooth muscle relaxation agents where both the compound of formula (A) is administered within 8 hours prior to sexual activity and the vascular smooth muscle relaxation agent is administered to the human within a sexually effective time period prior to sexual activity.
43. A method of inducing mating according to claim 42 where the vascular smooth muscle relaxation agents is selected from the group consisting of phosphodiesterase type 5 inhibitors, phosphodiesterase type 3 inhibitors, non-selective phosphodiesterase inhibitors, nitric oxide donor drugs, alpha type 1 adrenergic receptor antagonists, alpha type 2 adrenergic receptor antagonists, prostaglandin E1 receptor agonists (PGE1) and vasoactive intestinal polypeptide (VIP) agents.
44. A method of inducing mating according to claim 43 where the vascular smooth muscle relaxation agents is selected from the group consisting of sildenafil, ICOS-351, milrinone, papaverine, linsidomine, phentolamine, yohimbine, prostaglandin E1 (PGE1) and VIP.
45. A method of treating a sexual deficiency state in a human who has epilepsy, craniopharyngioma, hypogonadism or who has had a hysterectomy/oophorectomy, hysterectomy or oophorectomy which comprises administering a sexually therapeutically effective amount of a compound of the formula (A)



where

R₁, R₂ and R₃ are the same or different and are:

-H,

C₁-C₆ alkyl,

C₃-C₅ alkenyl,

5 C₃-C₅ alkynyl,

C₃-C₅ cycloalkyl,

C₄-C₁₀ cycloalkyl,

phenyl substituted C₁-C₆ alkyl,

10 -NR₁R₂ where R₁ and R₂ are cyclized with the attached nitrogen atom to
produce pyrrolidiyl, piperidiny, morphoniny, 4-methyl piperaziny or imidazolyl;

X is:

-H,

C₁-C₆ alkyl,

-F, -Cl, -Br, -I,

15 -OH,

C₁-C₆ alkoxy,

cyano,

carboxamide,

carboxyl,

20 (C₁-C₆ alkoxy)carbonyl,

A is:

CH,

CH₂,

CH-(halogen) where halogen is -F, -Cl, -Br, -I,

25 CHCH₃,

C=O,

C=S,

C-SCH₃,

C=NH,

30 C-NH₂,

C-NHCH₃,

C-NHCOOCH₃,

C-NHCN,

SO₂,

N;

B is:

CH₂,

CH,

5 CH-(halogen) where halogen is as defined above,

C=O,

N,

NH,

N-CH₃,

10 D is:

CH,

CH₂,

CH-(halogen) where halogen is as defined above,

C=O,

15 O,

N,

NH,

N-CH₃;

and n is 0 or 1, and where --- is a single or double bond, with the provisos:

20 (1) that when n is 0, and

A is CH₂, CH-(halogen) where halogen is as defined above, CHCH₃, C=O, C=S, C=NH, SO₂;

then D is CH₂, CH-(halogen) where halogen is as defined above, C=O, O, NH, N-CH₃;

25 (2) that when n is 0, and

A is CH, C-SCH₃, C-NH₂, C-NHCH₃, C-NHCOOCH₃, C-NHCN, N; then D is CH, N;

(3) that when n is 1, and

A is CH₂, CH-(halogen) where halogen is as defined above, CHCH₃, C=O, 30 C=S, C=NH, SO₂; and

B is CH₂, CH-(halogen) where halogen is as defined above, C=O, NH, N-CH₃; then

D is CH₂, C=O, O, NH, N-CH₃;

(4) that when n is 1, and

A is CH, C-SCH₃, C-NH₂, C-NHCH₃, C-NHCOOCH₃, C-NHCN, N; and

B is CH, N; then

D is CH₂, C=O, O, NH, N-CH₃;

(5) that when n is 1, and

5 A is CH₂, CHCH₃, C=O, C=S, C=NH, SO₂, and

B is CH, N; then

D is CH, N; and pharmaceutically acceptable salts thereof to the human.

46. A method of treating a sexual deficiency state according to claim 45 where the human is
10 a male.

47. A method of treating a sexual deficiency state according to claim 45 where the human is
a female.

15 48. A method of treating a sexual deficiency state according to claim 45 where the human
has epilepsy.

49. A method of treating a sexual deficiency state according to claim 45 where the human
has craniopharyngioma.

20 50. A method of treating a sexual deficiency state according to claim 45 where the human
has hypogonadism.

51. A method of treating a sexual deficiency state according to claim 45 where the human
25 has had a hysterectomy/oophorectomy.

52. A method of treating a sexual deficiency state according to claim 45 where the human
has had a hysterectomy.

30 53. A method of treating a sexual deficiency state according to claim 45 where the human
has a oophorectomy.

54. A method of treating a sexual deficiency state according to claim 45 where the

compound of formula (A) is administered orally, intra-nasally, buccally, intra-pulmonary, parenterally and rectally.

55. A method of treating a sexual deficiency state according to claim 54 where the
5 compound of formula (A) is administered orally, intra-nasally, buccally and intra-pulmonary.

56. A method of treating a sexual deficiency state according to claim 55 where the compound of formula (A) is administered orally.

10

57. A method of treating a sexual deficiency state according to claim 45 where the sexually therapeutically effective amount is from about 0.2 thru about 8 mg/person/dose.

58. A method of treating a sexual deficiency state according to claim 57 where the sexually
15 therapeutically effective amount is from about 0.5 thru about 5 mg/person/dose.

59. A method of treating a sexual deficiency state according to claim 58 where the sexually therapeutically effective amount is from about 1 thru about 3 mg/person/dose.

20 60. A method of treating a sexual deficiency state according to claim 45 where the compound of formula (A) is (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one.


61. A method of treating a sexual deficiency state according to claim 60 where the
25 compound of formula (A) is (5R)-5-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (Z)-2-butenedioate (1:1).

62. A method of treating a sexual deficiency state according to claim 45 where the pharmaceutically acceptable salt is selected from the group consisting of salts of the
30 following acids methanesulfonic, hydrochloric, hydrobromic, sulfuric, phosphoric, nitric, benzoic, citric, tartaric, fumaric, maleic, $\text{CH}_3-(\text{CH}_2)_n-\text{COOH}$ where n is 0 thru 4, $\text{HOOC}-(\text{CH}_2)_n-\text{COOH}$ where n is as defined above.

63. A method of treating a sexual deficiency state according to claim 45 where the compound of formula (A) is administered from about 10 minutes to about 8 hr prior to sexual activity.
- 5 64. A method of treating a sexual deficiency state according to claim 63 where the compound of formula (A) is administered from about 0.5 hr to about 1 hr prior to sexual activity.
65. A method of treating a sexual deficiency state according to claim 64 where the
10 compound of formula (A) is administered about 0.5 hr prior to sexual activity.
66. A method of treating a sexual deficiency state according to claim 45 where the mammal does not have Parkinson's disease.
- 15 67. A method of treating a sexual deficiency state according to claim 45 where the mammal does not experience postural hypotension.
68. A method of treating a sexual deficiency state according to claim 45 where the compound of formula (A) is used in combination with a sexually effective amount of one or
20 more vascular smooth muscle relaxation agents where both the compound of formula (A) is administered within 8 hours prior to sexual activity and the vascular smooth muscle relaxation agent is administered to the human within a sexually effective time period prior to sexual activity.
- 25 69. A method of treating a sexual deficiency state according to claim 68 where the vascular smooth muscle relaxation agents is selected from the group consisting of phosphodiesterase type 5 inhibitors, phosphodiesterase type 3 inhibitors, non-selective phosphodiesterase inhibitors, nitric oxide donor drugs, alpha type 1 adrenergic receptor antagonists, alpha type 2 adrenergic receptor antagonists, prostaglandin E1 receptor agonists (PGE1) and
30 vasoactive intestinal polypeptide (VIP) agents.
70. A method of treating a sexual deficiency state according to claim 69 where the vascular smooth muscle relaxation agents is selected from the group consisting of sildenafil, ICOS-

351, milrinone, papaverine, linsidomine, phentolamine, yohimbine, prostaglandin E1 (PGE1) and VIP.

71. A method of increasing sexual desire, interest or performance in a human who is
 5 desirous thereof which comprises administering a sexually useful effective amount of a compound of the formula (A)



where

R_1 , R_2 and R_3 are the same or different and are:

- 10 -H,
 C_1 - C_6 alkyl,
 C_3 - C_5 alkenyl,
 C_3 - C_5 alkynyl,
 C_3 - C_5 cycloalkyl,
 15 C_4 - C_{10} cycloalkyl,
 phenyl substituted C_1 - C_6 alkyl,
 - NR_1R_2 where R_1 and R_2 are cyclized with the attached nitrogen atom to
 produce pyrrolidyl, piperidinyl, morphoninyl, 4-methyl piperazinyl or imidazolyl;

X is:

- 20 -H,
 C_1 - C_6 alkyl,
 -F, -Cl, -Br, -I,
 -OH,
 C_1 - C_6 alkoxy,
 25 cyano,
 carboxamide,
 carboxyl,
 (C_1 - C_6 alkoxy)carbonyl,

A is:

CH,
 CH₂,
 CH-(halogen) where halogen is -F, -Cl, -Br, -I,
 CHCH₃,
 5 C=O,
 C=S,
 C-SCH₃,
 C=NH,
 C-NH₂,
 10 C-NHCH₃,
 C-NHCOOCH₃,
 C-NHCN,
 SO₂,
 N;

15 B is:

CH₂,
 CH,
 CH-(halogen) where halogen is as defined above,
 C=O,
 20 N,
 NH,
 N-CH₃,

D is:

CH,
 25 CH₂,
 CH-(halogen) where halogen is as defined above,
 C=O,
 O,
 N,
 30 NH,
 N-CH₃;

and n is 0 or 1, and where --- is a single or double bond, with the provisos:

(1) that when n is 0, and

A is CH₂, CH-(halogen) where halogen is as defined above, CHCH₃, C=O, C=S, C=NH, SO₂;

then D is CH₂, CH-(halogen) where halogen is as defined above, C=O, O, NH, N-CH₃;

5 (2) that when n is 0, and

A is CH, C-SCH₃, C-NH₂, C-NHCH₃, C-NHCOOCH₃, C-NHCN, N; then

D is CH, N;

(3) that when n is 1, and

10 A is CH₂, CH-(halogen) where halogen is as defined above, CHCH₃, C=O, C=S, C=NH, SO₂; and

B is CH₂, CH-(halogen) where halogen is as defined above, C=O, NH, N-CH₃; then

D is CH₂, C=O, O, NH, N-CH₃;

(4) that when n is 1, and

15 A is CH, C-SCH₃, C-NH₂, C-NHCH₃, C-NHCOOCH₃, C-NHCN, N; and

B is CH, N; then

D is CH₂, C=O, O, NH, N-CH₃;

(5) that when n is 1, and

20 A is CH₂, CHCH₃, C=O, C=S, C=NH, SO₂, and

B is CH, N; then

D is CH, N; and pharmaceutically acceptable salts thereof to the human.

72. A method of increasing sexual desire, interest or performance in a human who is desirous thereof according to claim 71 where the human is a male.

25

73. A method of increasing sexual desire, interest or performance in a human who is desirous thereof according to claim 71 where the human is a female.

74. A method of increasing sexual desire, interest or performance in a human who is
30 desirous thereof according to claim 71 where the compound of formula (A) is administered orally, intra-nasally, buccally, intra-pulmonary, parenterally and rectally.

75. A method of increasing sexual desire, interest or performance in a human who is desirous thereof according to claim 74 where the compound of formula (A) is administered orally, intra-nasally, buccally and intra-pulmonary.
- 5 76. A method of increasing sexual desire, interest or performance in a human who is desirous thereof according to claim 75 where the compound of formula (A) is administered orally.
77. A method of increasing sexual desire, interest or performance in a human who is
10 desirous thereof according to claim 71 where the sexually useful effective amount is from about 0.2 thru about 8 mg/person/dose.
78. A method of increasing sexual desire, interest or performance in a human who is desirous thereof according to claim 76 where the sexually useful effective amount is from
15 about 0.5 thru about 5 mg/person/dose.
79. A method of increasing sexual desire, interest or performance in a human who is desirous thereof according to claim 78 where the sexually therapeutic amount is from about 1 thru about 3 mg/person/dose.
20
80. A method of increasing sexual desire, interest or performance in a human who is desirous thereof according to claim 71 where the compound of formula (A) is (5R)-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one.
- 25 81. A method of increasing sexual desire, interest or performance in a human who is desirous thereof according to claim 80 where the compound of formula (A) is (5R)-5-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (Z)-2-butenedioate (1:1).
82. A method of increasing sexual desire, interest or performance in a human who is
30 desirous thereof according to claim 71 where the pharmaceutically acceptable salt is selected from the group consisting of salts of the following acids methanesulfonic, hydrochloric, hydrobromic, sulfuric, phosphoric, nitric, benzoic, citric, tartaric, fumaric, maleic, $\text{CH}_3-(\text{CH}_2)_n-\text{COOH}$ where n is 0 thru 4, $\text{HOOC}-(\text{CH}_2)_N-\text{COOH}$ where n is as defined above.

83. A method of increasing sexual desire, interest or performance in a human who is desirous according to claim 71 where the compound of formula (A) is administered from about 10 minutes to about 8 hr prior to sexual activity.

5

84. A method of increasing sexual desire, interest or performance in a human who is desirous according to claim 83 where the compound of formula (A) is administered from about 0.5 hr to about 1 hr prior to sexual activity.

10 85. A method of increasing sexual desire, interest or performance in a human who is desirous thereof according to claim 84 where the compound of formula (A) is administered about 0.5 hr prior to sexual activity.

15 86. A method of increasing sexual desire, interest or performance in a human who is desirous thereof according to claim 71 where the mammal does not have Parkinson's disease.

20 87. A method of increasing sexual desire, interest or performance in a human who is desirous thereof according to claim 71 where the mammal does not experience postural hypotension.

25 88. A method of increasing sexual desire, interest or performance in a human who is desirous thereof according to claim 71 where the compound of formula (A) is used in combination with a sexually effective amount of one or more vascular smooth muscle relaxation agents where both the compound of formula (A) is administered within 8 hours prior to sexual activity and the vascular smooth muscle relaxation agent is administered to the human within a sexually effective time period prior to sexual activity.

30 89. A method of increasing sexual desire, interest or performance in a human who is desirous thereof according to claim 88 where the vascular smooth muscle relaxation agents is selected from the group consisting of phosphodiesterase type 5 inhibitors, phosphodiesterase type 3 inhibitors, non-selective phosphodiesterase inhibitors, nitric oxide donor drugs, alpha type 1 adrenergic receptor antagonists, alpha type 2 adrenergic receptor

antagonists, prostaglandin E1 receptor agonists (PGE1) and vasoactive intestinal polypeptide (VIP) agents.

90. A method of increasing sexual desire, interest or performance in a human who is
5 desirous thereof according to claim 89 where the vascular smooth muscle relaxation agents is selected from the group consisting of sildenafil, ICOS-351, milrinone, papaverine, linsidomine, phentolamine, yohimbine, prostaglandin E1 (PGE1) and VIP.

91. (5R)-5-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinoline-2(1H)-thione and
10 pharmaceutically acceptable salts thereof.

92. A compound according to claim 91 which is (5R)-5-(Methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinoline-2(1H)-thione malate.

93. A method of treating sexual disturbances according to claim 1 where the compound of
15 formula (A) is (5R)-5-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinoline-2(1H)-thione.

94. A method of inducing mating according to claim 27 where the compound of formula
20 (A) is (5R)-5-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinoline-2(1H)-thione.

95. A method of treating a sexual deficiency state according to claim 45 where the
compound of formula (A) is (5R)-5-(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-
ij]quinoline-2(1H)-thione.

25

96. A method of increasing sexual desire, interest or performance in a human who is
desirous thereof according to claim 71 where the compound of formula (A) is (5R)-5-
(methylamino)-5,6-dihydro-4H-imidazo[4,5,1-ij]quinoline-2(1H)-thione.

30

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau




(43) International Publication Date
13 July 2000 (13.07.2000)

PCT

(10) International Publication Number
WO 00/40226 A3

- (51) International Patent Classification⁷: A61K 31/4745, 45/06, A61P 15/00
- (21) International Application Number: PCT/US99/27951
- (22) International Filing Date:
20 December 1999 (20.12.1999)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/114,840 6 January 1999 (06.01.1999) US
60/115,051 8 January 1999 (08.01.1999) US
60/115,922 14 January 1999 (14.01.1999) US
60/120,543 17 February 1999 (17.02.1999) US
- (71) Applicant (for all designated States except US): PHARMACIA & UPJOHN COMPANY [US/US]; 301 Henrietta Street, Kalamazoo, MI 49001 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): MEGLASSON, Martin, Durham [US/US]; 5337 Whippoorwill Drive, Kalamazoo, MI 49002 (US). McCALL, Robert, B. [US/US]; 413 Sunview, Kalamazoo, MI 49002 (US).
- (74) Agent: STEIN, Bruce; Intellectual Property Legal Services, Pharmacia & Upjohn Company, 301 Henrietta Street, Kalamazoo, MI 49001 (US).
- (81) Designated States (national): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— With international search report.
— Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.
- (88) Date of publication of the international search report:
1 February 2001
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHOD OF TREATING SEXUAL DISTURBANCES



(A)

(57) Abstract: The present invention is a method of treating sexual disturbances in humans and inducing mating in non-human mammals using the compounds of formula (A) in a dosage range where the sexually therapeutic amount is from about 0.2 thru 8 mg/person/dose and where the sexually mating amount is from about 0.003 thru 0.2 mg/kg/dose.

WO 00/40226 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/27951

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/4745 A61K45/06 A61P15/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data, MEDLINE, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 90 15058 A (UPJOHN CO) 13 December 1990 (1990-12-13)</p> <p>abstract page 2, line 14-24; claims 1-15,17; examples 1-137</p> <p>---</p> <p>-/--</p>	<p>1-15, 19-24, 27-35, 39-42, 45-59, 63-68, 71-79, 83-88</p>

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

8 document member of the same patent family

Date of the actual completion of the international search

12 October 2000

Date of mailing of the international search report

28/11/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

A. Jakobs

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 99/27951

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 95 04056 A (UPJOHN CO ;ROMERO ARTHUR G (US)) 9 February 1995 (1995-02-09) abstract page 2, line 4-17 page 17 ---	1-24, 27-42, 45-68, 71-88
E	US 6 018 046 A (KIKUCHI A ET AL) 25 January 2000 (2000-01-25) the whole document column 43, line 54 -column 44, line 8 ---	1-88
Y	DURHAM, ROBERT A. ET AL: "Effects of selective activation of dopamine D2 and D3 receptors on prolactin secretion and the activity of tuberoinfundibular dopamine neurons" EUR. J. PHARMACOL. (1997), 335(1), 37-42 , 1997, XP000951475 abstract page 39, column 1, paragraph 4 ---	1,3,4,6, 8,10-23, 45,47, 54-67
Y	HILDEBRANDT: "PSCHYREMBEL KLINISCHES WOERTERBUCH" , WALTER DE GRUYTER , BERLIN XP002149893 24017 page 1294, column 2, paragraph 11 -page 1295, column 1, paragraph 1 ---	1,3,4,6, 8,10-23, 45,47, 54-67
A	HEIER RICHARD F ET AL: "Synthesis and biological activity of (R)-5,6-dihydro-N,N-dimethyl-4H-imidazo(4,5,1-ij)quinolin-5-amine and its metabolites." JOURNAL OF MEDICINAL CHEMISTRY, vol. 40, no. 5, 1997, pages 639-646, XP002149892 ISSN: 0022-2623 the whole document -----	1-88

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Present claims 24,25,42,43,68,69,88,89 relate to a compound/therapeutic method defined (inter alia) by reference to the following parameter(s):

P1: a vascular smooth muscle relaxation agent.

P2: a phosphodiesterase type 3 inhibitor, a phosphodiesterase type 5 inhibitor, a non-selective phosphodiesterase inhibitor, nitric oxide drug donor, alpha type 1 adrenergic receptor antagonist, alpha type 2 adrenergic receptor antagonist, prostaglandin E1 receptor agonist, vasoactive intestinal polypeptide agents.

The use of these parameters in the present context is considered to lead to a lack of clarity within the meaning of Article 6 PCT. It is impossible to compare the parameters the applicant has chosen to employ with what is set out in the prior art. The lack of clarity is such as to render a meaningful complete search impossible. Consequently, the search has been restricted to the claimed uses of compounds of formula (A) alone or in combination with the compounds specifically mentioned in the claims and to the compound specified in claims 91-96.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/27951

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9015058 A	13-12-1990	AT 117688 T	15-02-1995
		AU 626427 B	30-07-1992
		AU 5743890 A	07-01-1991
		CA 2051697 A,C	10-12-1990
		DE 69016430 D	09-03-1995
		DE 69016430 T	01-06-1995
		DK 480939 T	10-07-1995
		EP 0480939 A	22-04-1992
		ES 2067744 T	01-04-1995
		FI 96310 B	29-02-1996
		FI 944704 A,B,	07-10-1994
		HK 65397 A	23-05-1997
		HU 210264 B	28-03-1995
		HU 9500660 A	28-11-1995
		JP 2955358 B	04-10-1999
		JP 4506071 T	22-10-1992
		KR 167346 B	15-01-1999
		NO 301421 B	27-10-1997
		RU 2023712 C	30-11-1994
		US 5436240 A	25-07-1995
		US 5273975 A	28-12-1993
WO 9504056 A	09-02-1995	AT 159943 T	15-11-1997
		AU 684808 B	08-01-1998
		AU 7246194 A	28-02-1995
		CA 2166700 A	09-02-1995
		CN 1128030 A,B	31-07-1996
		DE 69406678 D	11-12-1997
		DE 69406678 T	26-03-1998
		DK 724584 T	25-05-1998
		EP 0724584 A	07-08-1996
		ES 2108474 T	16-12-1997
		GR 3025482 T	27-02-1998
		JP 9500898 T	28-01-1997
		MX 9405676 A	31-01-1995
		NZ 269018 A	20-12-1996
		US 5652245 A	29-07-1997
US 6018046 A	25-01-2000	EP 0985671 A	15-03-2000
		WO 9745427 A	04-12-1997